

PHYSIOLOGICAL, BIOCHEMICAL AND MICROMORPHOLOGICAL CHANGES IN EPICUTICULAR WAXIN LEAVES OF CITRUS LEAVES (*Citrus aurantium* L.) INDUCED BY OZONE IN EGYPT

Ibrahim HASSAN

Department of Botany, Faculty of Science, Alexandria University, 21526 El Shatby, Alexandria. EGYPT

Email to i_hassan82@hotmail.com

Abstract

Macro- and micromorphological changes in epicuticular wax, the peroxidase activity, chlorophyll contents, photosynthetic rates (A), stomatal conductance (gs) and transpirational rates were investigated in the leaves of citrus plants (*Citrus aurantium* L.) grown in rural area un northern Egypt characterised by high levels of ambient ozone (O₃).

Exposure to ambient O₃ caused an increased rates of weathering of epicuticular wax, peroxidase activity, stomatal conductance and transpirational rate by 43, 40, 37 and 39%, respectively, while photosynthetic rates and chlorophyll contents were decreased by 35 and 26%, respectively.

Keywords: ambient ozone (O₃), Egypt, citrus leaves, epicuticular wax, photosynthetic rates, peroxidase activity.

INTRODUCTION

Decline of trees observed in Europe, the USA and other industrialised parts of the world has led to speculation on possible effect of atmospheric pollution [4].

Citrus trees (*Citrus aurantium* L.) are distributed in the Delta, Northern Egypt, the main agricultural land. However, a significant decline (defoliation and/or dieback) has been observed in recent years (Awad, person. Communication). Although several factors have been suggested to explain this phenomenon, the direct effect of oxidants was initially suspected [21], since the area of decline coincides well with the distribution of such air pollutants [20,22]. Soil acidification is also a possible factor, especially near tree bases. In addition, water deficiency may be a significant factor due to urbanisation [12, 20,42]

Nashimoto *et al* [29] reported a decline in cedar trees (*Cryptomeria japonica* D Don) in Japan due to oxidant pollutants, soil acidification and water stress [27 & 33]

Structural degradation of epicuticular wax has been attributed to the influence of O₃ [17, 35, 36 & 37], which may imply, that this degradation is coupled with a reduction of wax.

Increased leaching of ions has been reported to be another effect of O₃, which would lead to higher conductivity of leaves [17]

To the present knowledge, however, there has been little information about the physiological and functional changes in citrus trees in area of decline.

With regard to physiological changes, it has been suggested that oxidant air pollutants could affect the cuticle layer, especially the epicuticular wax, directly or indirectly [2, 3, 26, 35, 36 & 39]. In their studies Mengel *et al.*[28]; Barnes & Brown [2,3] ; Cape [9], reported that in *Picea abies* (L.) Karst, acid deposition and oxidant air pollutant accelerates melting and/or erosion of epicuticular

wax and change its chemical composition, thus increasing surface wettability [2, 7]. Since the cuticular layer plays an important role against water loss and pathogen infection, these changes may have serious effects on physiological functions of plants such as increase the cuticular transpiration rate [28, 35 & 36]

Plants tend to produce a large quantity of epicuticular wax under drought conditions, probably to compensate the lost water. Bondada *et al.* [6] reported that cotton plants (*Gossypium hirsutum* L.) grown under drought conditions showed increased amounts of epicuticular wax. These results were supported by results of Saneoka & Ogata [34], who reported similar results with forage crops.

Therefore, degradation of wax may cause serious water stress and sometimes lethal damage to plants [35, 36]

Amount of epicuticular wax may be affected by leaf age [8, 9 & 31] and by its morphology and chemical composition [16]. Nevertheless, structural degradation of epicuticular wax has been attributed to the influence of O₃ [37], which might imply that this degradation is coupled with a reduction of wax [2, 3 & 17]. However, several studies indicated that the properties of epicuticular wax may be affected not only by oxidant air pollution but also by natural environmental factors such as leaf age, morphology and composition of wax [2, 3, 8, 9, 10, 17, 31 & 36].

There is a lack of experimental evidence concerning the effects of O₃ on growth, physiology, biochemistry, and the micromorphology of epicuticular waxes of Egyptian trees.

The present study was undertaken to fill the above mentioned gap of knowledge and to clarify the alteration in epicuticular wax, photosynthetic rates, stomatal conductance and peroxidase levels (as biochemical marker in response to O₃ stress) in leaves of citrus (*Citrus aurantium* L.).

MATERIALS AND METHODS

Sampling of citrus leaves under Egyptian field conditions

Leaves of *Citrus spp.* were collected from a rural site in northern Egypt (Abbis village). The description of the site and topography are discussed elsewhere [22]. Sampling of leaves was carried out from 5 years old trees and was carried out from October to April for three consecutive years (2001 – 2003). Controlled samples were collected in parallel with samples from trees grown in unpolluted glasshouse.

The leaves were collected in icebox and transferred to the laboratory and kept at 5 °C overnight until analysis in the following day.

There was no any microbial infection or morphological changes in the wax under these conditions [32, 35 & 36]. Moreover, under these conditions, leaves remained fresh without any water loss.

During transportation and keeping of leaves, care was taken not to abrade the wax.

Photosynthetic rates (A) and stomatal conductance (gs) measurements

A and gs were measured once a week, regularly on the same leaf throughout the entire course of the experiment over the three successive years using LICOR-1600 (Licor, Nicolin, USA)

Extraction of Epicuticular wax

Epicuticular wax was extracted according to [8] and the modified method [35 & 36] Forty grams of fumigated and non-fumigated leaves was taken from different branches on the trees. Leaves were washed with deionised water for about 5 min (the washed leaves showed no loss in the wax due to washing). The washed leaves were dried in an oven at 39 °C for 45 min and allowed to dry on a laboratory bench at room temperature (25 – 30 °C). The weight of prepared leaves was regarded as fresh mass, since the water content of leaves remains constant throughout the whole process, including storage and washing [35 & 36]. Three grams of prepared leaves was shaken for 25 sec with 30 ml chloroform to extract epicuticular wax. The wax extracts were then filtered through quartz wool and a filter paper (Watmann No. 3) to remove aerosols adsorbed on the leaf surface. Six batches were prepared initially for each set of fumigated and non-fumigated leaves, and two of them were combined to make triplicate composite extract. The chloroform was then evaporated and

the residual wax was determined gravimetrically (Barnes, J.D., Pers. Communication). The amount of wax was expressed on a fresh mass (mg/g fresh leaf weight).

The wax samples were refrigerated using a filter Cartridge (Millipore, pore size 0.5 μm) and stored in liquid nitrogen (-80°C) for analysis of chemical composition.

Measurement of cuticular transpiration

Twenty leaves rinsed with distilled water and dried gently against a paper towel at a room temperature. The cut part of leaves was covered with grease to avoid excessive water loss.

Leaves were weighed and then stored in a desiccator at ($25 - 30^{\circ}\text{C}$) in the laboratory. Leaves were weighed daily at 12:00 h until the weight was constant (after 7 days) which indicates cuticular transpiration [28].

Determination of elemental composition of epicuticular wax

The C, N and O contents were determined according to Sase *et al.* (1998) with an elemental analyser (Fisons, EA 1110). Sulfanilamide ($\text{C}_6\text{H}_8\text{O}_2\text{N}_2\text{S}$, Fisons) was used as a standard sample for N, 1-Docosanol ($\text{C}_{22}\text{H}_{46}\text{O}$, Fluka Chemie) was used for O and benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$, Fisons) for N.

The analyses were repeated twice for each sample.

Statistical analysis

One-way ANOVA was applied to log-transformed data. There was no covariate used except for photosynthetic rates and stomatal conductance, where photosynthetic flux density (PPFD) was used as a covariate. Significant difference between means tested by LSD (STATGRAPHICS Statistical Package).

RESULTS

The mean concentrations of major air pollutants (O_3 , SO_2 and NO_x) through the growing seasons are shown in table 1.

Pollutant				
Year	Experiment	O_3 (8h d^{-1})	SO_2 (6h d^{-1})	NO_x (6h d^{-1})
2001	NF	71	10	7
	FA	11	3	3
2002	NF	79	8.7	6
	FA	10	3	2
2003	NF	91	7	5
	FA	11	3	2

Table 1. Mean concentrations (nl l^{-1}) of major pollutants during the growing seasons at Abbis village in the charcoal-filtered (FA) and non-filtered (NF) OTCs.

Air quality at the experimental site was characterised by very low concentrations of SO_2 and NO_x , where the mean 6-h concentrations of these gases through the four growing seasons were 8.5 and 6.4 nl l^{-1} , respectively (table 1). The mean 8-hour concentrations of O_3 through the growing seasons (2001–2003) averaged 80 nl l^{-1} (table 1).

Table 2 shows the amount of wax collected from leaves of citrus plants grown in filtered and non-filtered air. The amount of epicuticular wax (mg/g dry leaves) was found to decrease year by year

due to exposure to unfiltered air: it was 35, 42 and 51% reduction in 2001, 2002 and 2003, respectively. Moreover, there was a strong positive correlation between the loss of wax and the ambient O₃ levels in the study area ($R^2 = 1.10$, data not shown).

Experiment			
tment	0/ 2000 – 4/2001)	0/2001 – 4/2002)	0/2002 – 4/2003)
	19.3	20.1	19.9
	12.6 (-35%)	11.7 (-42%)	9.8 (-51%)

Table 2. Amount of wax (mg/g fresh leaf); CF = Charcoal-filtered air, NF = non-filtered air; Figures between parthenses are percentage difference from the control.

The chlorophyll content (mg/g) of leaves was shown in table 3. It was indicated that chlorophyll contents of plants grown in non-filtered air was significantly lower than that collected from plants grown in filtered air (averaged 35% for the three seasons). Moreover, the highest percentage reduction in chlorophyll was recorded to be -42% in 2003 and this was also significantly correlated to O₃ levels recorded in the region ($R^2 = 0.97$).

Experiment			
tment	0/ 2000 – 4/2001)	0/2001 – 4/2002)	0/2002 – 4/2003)
	8.2	7.4	7.3
	5.8 (-29%)	5.1 (-31%)	4.4 (-42%)

Table 3. Chlorophyll content (mg/g fresh leaf); Legends as table 1.

Table 4. shows that C,N and O contents of epicuticular wax. Parallel with the decreased in chlorophyll contents of leaves and the loss of wax, the concentrations of C,N and O contents were significantly lowered in non-filtered plants.. The lowest percentage reduction was in C content (ca. 20%), while that of N and O contents was 56% each (averaged between the three experimental seasons).

Experiment								
	0/ 2000 – 4/2001)			0/2001 – 4/2002)			0/2002 – 4/2003)	
tment								
	4						3	
	5			8			7	

Table 4. Elemental composition of epicuticular wax; Legends as table 1.

To evaluate the effect of wax reduction on plant physiology, cuticular transpiration rate was examined (Table 5). Exposure to non-filtered air caused an increase in cuticular transpiration, and the highest percentage increase (ca.64%) was found in 2003 and this was correlated to the increase in O₃ levels ($R^2 = 0.85$) and the loss in wax ($R^2 = 0.98$).

Experiment			
tment	0/ 2000 – 4/2001)	0/2001 – 4/2002)	0/2002 – 4/2003)
	0.45	0.53	0.61
	0.61	0.82	1.36

Table 5. Cuticular transpiration rate (%h); Legends as table 1.

Table 6. Shows the there was an increase in peroxidase activity in leaves exposed to non-filtered air (averaged ca. 40%) compared to that extracted from plants grown in filtered air.

Experiment			
tment	0/ 2000 – 4/2001)	0/2001 – 4/2002)	0/2002 – 4/2003)
	5.0	8.4	4.5
	5.6	8.9	9.4

Table 6. Peroxidase activity ($\mu\text{mol g}^{-1}$ fresh weight) in leaves of *Citrus aurantium* L.; Legends as table 1.

Figure 1 a. Shows that photosynthetic rate (A) of plants grown in non-filtered air was decreased by about 26%. On the other hand, stomatal conductance (g_s) was found to be increased by about 37% (Fig 1b).

DISCUSSION

The concentrations of O_3 recorded in the present study further a support for previous results [22 & 42] that suggested that it is unlikely to have SO_2 and NO_x in a rural site in Egypt, which has been supported recently [20]. Moreover, the results of the present investigation are in agreement with the results of Anjea *et al.* [1] who reported relatively high levels of O_3 in a rural site in the USA and Schenone & Lorenzini [38] who had similar results in Italy.

It seems very reasonable to conclude that ambient O_3 played a major role in causing the observed alterations in morpho-functional parameters examined here (epicuticular wax and peroxidase) in plants exposed to non-filtered air.

The observed changes in epicuticular wax structure are in agreement with the effects of pollutants until now described for coniferous species [25, 35, 36 & 37], as there are only very few studies investigating the effects of air pollutants on degradation of waxes on broad-leaved plants [6 & 26]. Manes *et al* [26] found a serious alteration of the epicuticular waxes in *Nicoaiana tabacum* leaves collected from urban areas, where there was a greater accumulation of particulate matter and epicuticular waxes formed an amorphous fused layer.

In general, waxes decrease with age [7, 8, 31, 35 & 36]. This reduction occurs and is sometimes accelerated by factors such as wind [18, 19 & 41] as well as by air pollution [2, 23, 24 & 40].

It was suggested that air pollution contribute to the erosion of wax. Therefore, the relatively low amount of epicuticular wax in leaves collected from non-filtered plants. Moreover, the amount of epicuticular waxes collected in 2003 was lower than that collected in the other two seasons, this may have resulted from aging of leaves, affected mainly by higher O_3 levels recorded in 2003.

The increased stomatal conductance (reduced electrical conductivity) was suggested to be resulted from impeded uptake or plant-internal distribution of the ions, especially there was loss in elemental composition of leaves, especially in N content. However, the decreased contents of measured elements cannot alone cause the total reduction. It is also possible that fewer organic substances had leached into the diffusate especially there was degradation in epicuticular wax [26].

The negative correlation ($R^2 = -0.98$) between cuticular transpiration and the amount of wax, could provide a strong evidence for the potential stress due to O_3 in the region, which may have altered the characteristics of epicuticular wax in citrus tress grown in non-filtered conditions and this is in agreement with the results of O'Toole *et al.* [30]; Bengston *et al.* [5] and Jordon *et al.* [24] on rice, oat and sorghum, respectively. These authors reported that cuticular transpiration rates increased after removal of wax. It is suggested that the amount of wax plays a key role in regulating the cuticular transpiration rate in citrus plants. Moreover, the increased stomatal conductance (g_s) may cause an increase in cuticular transpiration especially there was stomatal malfunction observed (data not shown), this is in agreement with the results that reported an increase in g_s of radish leaves after exposure to 89 nl l^{-1} [20] and they attributed this increase in g_s to destruction of epidermal cells and stomatal malfunction. This phenomenon

has been suggested to cause incomplete stomatal closure of stomata, resulting in water loss [11, 13, 14 & 35].

A rise in peroxidase activity due to ozone stress, as occurs in leaf senescence [5, 10 & 26], would represent an induced protective reaction. This enzymatic system plays an important role in the maintenance of an adequate redox potential in leaf cells and may protect the cell membrane from active oxidant arising from exogenous sources [15]. Although peroxidase test has proved to be a valid marker of acid stress, more work is needed to evaluate this test as marker of a stress attributable to atmospheric pollution.

In Egypt, the temperature is increasing, while the precipitation has decrease year after year and thus the atmosphere has shifted towards dry conditions. Therefore, it is likely that citrus trees suffer significant water stress, not only because of enhanced transpiration through the combined effect of degraded wax layer and partial malfunctioning of stomata, but also because the recent dry conditions. This stress may prove lethal at times, especially there are some unexplained symptoms appeared on trees in the field such as dieoff and/or defoliation of treetop, even though the lower part of trees may be apparently healthy.

Nevertheless, the parameters examined in the present study showed that the plants respond to O₃ with a pattern similar to that described for senescence. Therefore, the action stress may be interpreted as an accelerative process of aging.

In general, the results of the present investigation are important not only as basic information on the physiological properties of Citrus plants, but also for future evaluation of the effect of anthropogenic environmental factors on plants.

In conclusion, O₃ accelerate the natural aging of wax and increased frequency of stomata occluded by an amorphous plug, which would be expected to have far reaching physiological consequences in vivo, such as increased number of diseased and dying trees in Egypt.

More work, however, is urgently needed to identify the effects of season and the relationships between the impact of O₃ on tree species growing in natural sites, which differ in exposure and average concentrations of major gaseous pollutants.

ACKNOWLEDGEMENTS:

The present work is partially funded with a grant from the British Council (Higher Educational Link, DFID)

I would like to express my deepest thanks to Dr. J.D. Barnes, Newcastle University, UK for his help in extraction and determination of epicuticular wax and Prof. J.N.B. Bell, Imperial College, London for hosting me at his laboratory during preparing and writing up the present work.

REFERENCES

- [1] Anjea, V.P., Yoder, G.T. & PalArua, S.1992. O₃ in the urban southeastern United States. *Environ. Pollut.* **75**, 39 – 44.
- [2] Barnes, J. D. and Brown, G.M.1990. The influence of O₃ and acid mist on the amount and wettability of the surface waxes in Norway spruce (*Picea abies* L. Karst). *New Phytol.* **114**: 531 – 535.
- [3] Barnes, J.D., Eamus, D., Davison, A.W., Ro-Poulsen, H. and Mortensen, L. 1990. Persistent effects of O₃ on needle water loss and wettability in Norway spruce. *Environ. Pollut.* **63**, 345 – 363.
- [4] Bell, J.N.B. 1986. Effects of acid deposition on crops and forest. *Experientia* **42**: 363 – 371.
- [5] Birecka, H., Chaskes, M.J. and Goldstein, J. 1979. Peroxidase and senescence. *J. Exp. Bot.* **30**(116): 565 – 573.
- [6] Bondada, B.R., Oosterhuis, D.M., Murphy, J.B., and Kim, K.S. 1996. Effects of water stress on the epicuticular wax composition and ultrastructure of cotton (*Gossypium hirsutum* L.) leaf, bract and ball. *Environ. Exp. Bot.* **36**: 61 69.
- [7] Cape, J.N. 1983. Contact angles of water droplets on needles of Scots pine (*Pinus sylvestris*) growing in polluted atmospheres. *New Phytol.* **93**: 293 – 299.

- [8] Cape, J.N. 1986. Effects of air pollution on the chemistry of surface waxes of Scots pine. *Water Air and Soil Pollution* **31**: 393 – 399.
- [9] Cape, J.N., Paterson, I.S. and Wolfenden, J. 1989. Regional variation in surface properties of Norway spruce and Scots pine needles in relation to forest decline. *Environ. Poll.* **58**: 325 – 342.
- [10] Castillo, F.J. 1986. Extracellular peroxidase as markers of stress. In (Greppin, H., Penel, Cand Th. Gaspar, Eds.) *Molecular and Physiological Aspects of Plant Peroxidase*. University of Geneva, Switzerland, pp. 10 – 21.
- [11] Crossely, A. and Fowler, D. 1986. The weathering of Scots pine epicuticular wax in polluted and clean air. *New Phytol.* **103**: 207 – 218.
- [12] El-Khatib, A.A. 2003. The response of some Egyptian plants to O₃ and their use as biomointors. *Environ. Pollut.* **124**: 419 – 428.
- [13] Eveling, D.W. 1986. Scanning electron microscopy of damage by dust deposits to leaves and petals. *Bot. Gaz.* **147**: 159 – 165.
- [14] Eveling, D.W. and Bataillé, A. 1984. The effects of deposits of small particles on the resistance of leaves and petals to water loss. *Environ. Pollut.* **36**: 229 – 238.
- [15] Fridovich, I. 1978. The biology of oxygen radicals. *Sci.*, **201**: 875 – 880.
- [16] Günthardt-Goerg, M.S. 1986. Epicuticular wax of needles of *Pinus cembra*, *Pinus sylvestris* and *Picea abies*. *Eur. J. Forset Path.*, **16**: 400 – 418.
- [17] Günthardt-Goerg, M.S. and Keller, T. 1988. Cuticular features and conductivity of needle diffusate in spruce. In. (Cape, J.N. & Mathy, P. Eds.) *Scientific Basis of Forest Decline Symptomatology*. Commission of European Communities, Belgium, pp316 – 322.
- [18] Hadley, J.L. and Smith, W.K. 1983. Influence of wind exposures on needles desiccation and mortality for timberline conifers in Wyoming, USA. *Arct. Alp. Res.* **15**: 127 – 135.
- [19] Hadley, J.L. and Smith, W.K. 1986. Wind effects on needles of timberline conifers: seasonal influence on mortality. *Ecology*, **67**: 12 – 19.
- [20] Hassan, I.A. 1999. Air pollution in Alexandria region, Egypt. I. An investigation of air quality. *Int. J. Environ. Educ. & Inform.* **18**(1): 67 - 78.
- [21] Hassan, I.A., Ashmore, M.R. & Bell, J.N.B. 1994. Effects of O₃ on the stomatal behaviour of Egyptian verities of radish (*Raphanus sativus* L. cv. Baladey) and turnip (*Brassica rapa* L. cv. Sultani). *New Phytol.* **128**, 243-49.
- [22] Hassan, I.A., Ashmore, M.R. & Bell, J.N.B. 1995. Effect of O₃ on radish and turnip under Egyptian field conditions. *Environ. Pollut.* **89**, 107-14.
- [23] Huttunen, S. 1994. Effects of air pollutants on epicuticular wax structure. In. *Air Pollutants and the Leaf Cuticle*. Springer-Verlag, Berlin, pp. 81 –69.
- [24] Jordon, W.R., Shouse, P.J. Blum, A. Miller, F.R., and Monk, R.L. 1984. Environmental physiology of sorghum. II. Epicuticular wax load and cuticular transpiration. *Crop Sci*, **24**: 1168 – 1173.
- [25] Karhu, M. and Huttunen, S. 1986. Erosion effects of air pollution of needle surface. *Water Air & Soil Pollut.*, **31**: 417 – 424.
- [26] Manes, F., Alteri, A., Anelini, F., Burno, M., Cortillo, L and Federico, R. 1988. Micromorphological and biochemical changes in *Pinus Pinea* L. and *Nicotiana tabacum* L. in relation to atmospheric pollutants. In. (Cape, J.N. & Mathy, P. Eds.) *Scientific Basis of Forest Decline Symptomatology*. Commission of European Communities, Belgium, pp 342 - 353
- [27] Matsumoto, Y., Maruyama, Y. and Morikawa,, Y. 1992. Some aspects of water relations on large *Cryptomeria japonica* leaves D. Don trees and climatic changes on the kanto plains in Japan in relation to forest decline. *Jap. J. For. Environ.* **34**: 2 –13.
- [28] Mengle, K., Hogrebe, A. and Esh, A. 1989. Effects of acidic fog on needle surface and water relations of *Picea abies*. *Physiol. Plant.*, **75**: 201 - 207.
- [29] Nashimoto, M., Takahashi, K., and Ashihara, S. 1993. Comparison of decline of Japanese cedar (*Cryptomeria japonica* D. Don) stands with soil chemical properties in the Kanto-Koshin Distict. *Environ. Sci.* **6**: 121 – 130.

- [30] O'Toole, J.C., Cruz, R.T. and Seiber, J.N. 1979. Epicuticular wax and cuticular resistance in rice. *Physiol. Plant.*, **47**: 239 – 244.
- [31] Percy, K.E., Jagles, R., Marden, S., McLaughlin, C.K., and Carlsil, J. 1993. Quantity, chemistry and wettability of epicuticular waxes on needles of red spruce along a fog-acidity gradient. *Can. J. For. Res.* **23**: 1472 – 1479.
- [32] Percy, K.E., McQuattie, C. J and Rebbeck, J.A. 1994. Effects of air pollution on epicuticular waxchemical composition. In *Air Pollutants and The Leaf Cuticle*. NATO ASI Series Vol. G 36. Springer_Verlag, Berlin, pp. 67 – 79.
- [33] Sakata, M. 1996. Evaluation of possible causes for the decline of Japanese cedar (*Cryptomeria japonica*) based on elemental composition and $\delta^{13}\text{C}$ of needles. *Environ. Sci. Technol.*, **30**: 2376 – 2381.
- [34] Saneoka, H. and Ogata, S. 1987. Relationship between water use efficiency and cuticular wax deposition in warm season forage crops grown under deficit conditions. *Soil Sci. Plant Nutr.* **33**: 439 – 448.
- [35] Sase, H., Takamatsu, T. and Yoshida, T. 1998a. Variation in amount and elemental composition of epicuticular wax in Japanese cedar (*Cryptomeria japonica*) leaves associated with natural environmental factors. *Can. J. Forest Res.* **28**(1): 87 – 97.
- [36] Sase, H., Takamatsu, T., Yoshida, T. and Inubushi, K. 1998b. Changes in properties of epicuticular wax and the related water loss in Japanese cedar (*Cryptomeria japonica*) affected by anthropogenic environmental factors. *Can. J. Forest Res.* **28**(4): 546 – 556.
- [37] Sauter, J.J. and Voss, J.V. 1986. SEM observations on the structural degradation of epistomatal waxes in *Picea abies* (L.) Karst. And its possible role in the "Fichtensterben". *Eur. J. For. Path.*, **16**: 408 – 423.
- [38] Schenone, G. & Lorenzini, G. 1992. Effects of regional air pollution on crops in Italy. *Agric. Ecosyst. Environ.* **38**,55-66
- [39] Turunen, M., and Huttunen, S., 1990. A review of the response of epicuticular wax of conifer needles to air pollution. *J. Environ. Qual.* **19**: 35 – 45.
- [40] Turunen, M., Huttunen, S., Percy, K.E., McLaughlin, C.K. and Lamppu, J. 1997. Epicuticular wax of subarctic Scots pine needles: response to sulphur and heavy metals deposition. *New Phytol.* **135**: 501 – 515.
- [41] Wilson, J. 1984. Microscopic features of wind damage to leaves of *Acer pseudoplatanus* L. *Ann. Bot.*, **53**: 73 – 82.
- [42] WHO/UNEP 1993. *Air Pollution in Megacities of The World*. World Health Organisation, United Nations Environment Programme. Blackwell, Oxford.